



**E.F.B.E.
1954**

FREE FULL TEXT

This is a summary plus of a paper published in Hellenic Journal of Physical Education and Sport Science. You can cite this article as follows:

I. Giannopoulou, H. Kilduff, M. Kingsley, I. Vogiatzis, C. Roussos , G. Moschandreas, A. Kafatos, J. McEneny, A. L. Skaltsounis, I. P. Pitsiladis (2010) Effects of olive oil phenols on exercise-induced oxidative stress markers in elite cyclists after a 3-week training program. *Hellenic J Phys Educ & Sport Sci*, 2010, (30) 2: 163 – 176

EFFECTS OF OLIVE OIL PHENOLS ON EXERCISE-INDUCED OXIDATIVE STRESS MARKERS IN ELITE CYCLISTS AFTER A 3-WEEK TRAINING PROGRAM*

I. Giannopoulou ¹, H. Kilduff ², M. Kingsley ², I. Vogiatzis ¹, C. Roussos ¹, G. Moschandreas ³, A. Kafatos ³, J. McEneny ⁴, A. L. Skaltsounis ⁵, I. P. Pitsiladis ⁶

¹ Thorax Institute, School of Medicine, Athens, Greece

² Department of Sports Science, University of Wales Swansea, Swansea, SA2 8PP, United Kingdom

³ Epidemiology and Social Health and dietetics clinic, Department of Social Medicine, Faculty of Medicine, University of Crete, Heraklion, Crete, Greece

⁴ Nutrition and Metabolism Group, The Queen's University Belfast, Grosvenor Road, Belfast, BT12 6BJ, Northern Ireland

⁵ Department of Pharmacy, University of Athens, Athens, Greece

⁶ Institute of Biomedical & Life Sciences, University of Glasgow, Glasgow, G12 8QQ, United Kingdom



Abstract

The purpose of the present study was to assess the effects of high and low phenol-enriched olive oil on markers of oxidation, plasma susceptibility to oxidation and antioxidant status of elite athletes. Fifteen elite level cyclists took part in a 3-week intervention periods with high and low phenolic content olive oil that were separated by a washout period of 2 weeks. During the intervention periods, 30 grams of olive oil were provided to each athlete per day. A two-way repeated measures analysis of variance showed that both olive oil phenol supplements resulted in significantly higher concentrations of Vitamin C ($P=0.001$). The lipid hydroperoxide concentration increased only after supplementation with the low phenol olive oil ($P=0.03$). No significant differences were found in any of the other measured antioxidant markers as a result of the olive oil phenols. These findings demonstrate a lack of a protective effect of olive oil phenols on the oxidative damage induced by strenuous exercise in elite athletes. It is possible that the background Mediterranean diet as well as the high fitness level of these elite athletes did not allow for an additional antioxidant effect to take place with the olive oil phenol supplementation. Future research is needed to further investigate this issue.

Key words: phenol, olive oil, oxidative stress, exercise, cycling

*An extended Summary Plus English version is freely available at www.hellenicjsport.com

Introduction

It is well known that strenuous high-intensity exercise can lead to disturbances in antioxidant homeostasis by elevating the concentration of free radicals and other reactive oxygen species (ROS) (Urso & Clarkson, 2003). ROS pose a serious threat to the cellular antioxidant defense system and has detrimental effects on exercise performance (Powers et al, 2004). To date, there is limited evidence that dietary supplementation with antioxidants improves physical performance (Ji, 1999; Powers et al, 2004; Urso & Clarkson, 2003). How olive oil, a food rich in antioxidant compounds, particularly due to its high content of phenols, affects the oxidative stress of elite athletes is not known. The purpose of the present study was to assess the effects of high and low phenol-enriched olive oil on markers of oxidation, plasma susceptibility to oxidation and antioxidant status of elite athletes. It was hypothesized that supplementation with high phenol-enriched olive oil would provide an enhanced antioxidant defense mechanism during strenuous exercise compared to low phenol-enriched olive oil supplementation.

Methods

Fifteen elite level cyclists with a maximal oxygen uptake of 62.0 ± 6.0 ml·kg⁻¹·min⁻¹ were recruited for the study. A double-blind, randomized crossover study was undertaken. The experimental study consisted of two 3-week intervention periods with high and low phenolic content olive oil that were separated by a washout period of 2 weeks. The enriched olive oil contained 30 mg of oleuropein in 30 mL of oil. During the intervention periods, 30 grams of olive oil (approximately 2 tablespoons) were provided to each athlete per day. The athletes were blind to the type of olive oil given.

Training during the two intervention periods comprised of prolonged endurance cycling interspersed with at least two high-intensity interval-training sessions per week. Blood antioxidant and oxidative stress markers were determined from two 10 ml samples collected after an overnight fast and after the ingestion of 30 grams of either low or high phenol-enriched olive oil followed by a standardized high-intensity interval-training session at the end of each 3-week intervention period. A two-way repeated measures analysis of variance was applied to all biochemical variables with SPSS for Windows, version 11 (SPSS Inc., IL). All values are presented as mean \pm SD.

Results

Both olive oil phenol supplements resulted in significantly higher concentrations of Vitamin C ($P=0.001$). The lipid hydroperoxide concentration increased only after supplementation with the low phenol olive oil ($P=0.03$). No significant differences were found in any of the other measured antioxidant markers as a result of the olive oil phenols

Table 1. Oxidative and antioxidant markers concentration pre and post high and low phenol interventions (n=15)

Dependent Variable	Low phenol		Hi phenol	
	Pre	Post	Pre	Post
LDL lag time (minutes)	65.6 ± 11.8	64.2 ± 10.2	69.9 ± 11.4	65.5 ± 9.3
Lipid Hydroperoxides ($\mu\text{mol}\cdot\text{L}^{-1}$)	0.80 ± 0.3	1.10 ± 0.6*	1.00 ± 0.5	1.15 ± 0.3
Vitamin C ($\mu\text{mol}\cdot\text{L}^{-1}$)	27.80 ± 17.2	64.13 ± 30.0**	37.81 ± 15.7	64.61 ± 22.4**

* P<0.05 from pre-intervention, **P<0.01 from pre-intervention

Discussion

This investigation is one of the first attempts to examine the effects of olive oil-derived phenols on oxidative damage induced by strenuous exercise in humans. In animals as well as in humans, administration of olive oil phenols results in reduced oxidative damage during resting conditions (Scaccini et al, 1992; Vissers et al, 2004). In the present study, strenuous exercise in combination with administration of the low olive oil phenol supplementation does not appear to produce clinically significant improvements in the oxidative state of the elite athletes. Specifically, exercise resulted in a significant elevation (approximately two-fold) in Vitamin C irrespective of the supplementation with the olive oil phenols. Our findings are in accordance with previous studies showing immediate increases in plasma Vitamin C levels as a result of a strenuous exercise session (Gleeson et al, 1987).

Only small changes in oxidative damage, such as suppression with the high phenol supplementation of lipid hydroperoxide concentration, a known consequence of free radical and ROS generation, as well as a lack of change in LDL lag time, also an indicator of plasma lipid susceptibility to oxidation were found. Such findings could possibly indicate a protective effect of the olive oil phenols on the oxidative damage induced by strenuous exercise, however more research is needed to establish such a phenomenon.

Conclusion

The findings of this study demonstrate a lack of a protective effect of olive oil phenols on the oxidative damage induced by strenuous exercise in elite athletes. It is possible that the background Mediterranean diet as well as the high fitness level of these elite athletes did not allow for an additional antioxidant effect to take place with the olive oil phenol supplementation. Since this is the first study to examine the effects of olive oil phenols on exercise-induced oxidative stress and as there is considerable lack of research on the antioxidant properties of olive oil phenols in humans, let alone athletes, no conclusive statements can be made at this point. Future research is needed to further investigate this issue.

References*

- Catignani, G.L. & Bieri, J.G. (1983): Simultaneous determination of retinol and alpha-tocopherol in serum or plasma by liquid chromatography. Clinical Chemistry, 29, 708-712.
- Clarkson, P.M. & Thompson, H.S. (2000): Antioxidants: what role do they play in physical activity and health? American Journal of Clinical Nutrition, 72, 637S-646S.
- Duthie, G.G., Robertson, J.D., Maughan, R.J., Morrice, P.C. (1990). Blood antioxidant status and erythrocyte lipid peroxidation following distance running. Archives of Biochemistry and Biophysics, 82, 78-83.
- Gleeson, M., Robertson, J.D., Maughan, R.J. (1987). Influence of exercise on ascorbic acid status in man. Clinical Science, 73, 501-505.
- Hochstein, P. & Ernster, L. (1963): ADP-activated lipid peroxidation coupled on TPNH oxidase system of microsomes. Biochemical and Biophysical Research Communications, 12, 388-394.
- Ji, L. (1999): Antioxidants and oxidative stress in exercise. The Society for Experimental Biology and Medicine, 222, 283-292.
- Masella, R., Vari, R., D'Archivio, M., Di Benedetto, R., Matarrese, P., Malo, W., Scazzocchio, B., Giovannini, C. (2004). Extra virgin olive oil biophenols inhibit cell-mediated oxidation of LDL by increasing the mRNA transcription of glutathione-related enzymes. Journal of Nutrition, 134, 785-791.
- Marrugat, J., Covas, M.I., Schroder, H., Miro-Casas, E., Gimeno, E., Lopez-Sabatar, M.C., De la Torre, R., Farre, M.; SOLOS Investigators (2004): Effects of differing phenolic content in dietary olive oils on lipids and lipid oxidation-a randomized controlled trial. European Journal of Nutrition, 43, 140-147.
- McEneny, J., Trimble, E.R., Young, I.S. (1998): A simple method for assessing copper-mediated oxidation of very-low-density lipoprotein isolated by rapid ultracentrifugation. Annals of Clinical Biochemistry, 35, 504-514.
- McDowell, I.F.W., McEneny, J., Trimble, E.R. (1995) A rapid method for measurement of the susceptibility to oxidation of low-density lipoprotein. Annals of Clinical Biochemistry, 32: 167-174
- Peake, J. (2003): Vitamin C: effects of exercise and requirements with training. International Journal of Sport Nutrition and Exercise Metabolism, 13, 125-151.
- Robertson, J.D., Maughan, R.J., Duthie, G.G., Morrice, P.C. (1991): Increased blood antioxidant systems of runners in response to training load. Clinical Science, 80, 611-618.
- Ross, D. & Moldeus, P. (1991): Antioxidant defense systems and oxidative stress. In: Vigo-Pelfrey, C., ed. Membrane lipid oxidation. Boca Raton, FL: CRC Press; 151-170.
- Sacheck, J.M. & Blumberg, J.B. (2001): Role of Vitamin E and oxidative stress in exercise. Nutrition, 17, 809-814.
- Scaccini, S., Nardini, M., D'Aquino, M., Gentili, V., Di Felice, M. & Tomassi, G. (1992): Effect of dietary oils on lipid peroxidation and on antioxidant parameters of rat plasma and lipoprotein fractions. Journal of Lipid Research, 33, 627-633.
- Thurnham, D.I., Smith, E., Flora, P.S. (1988): Concurrent liquid-chromatographic assay of retinol, alpha-tocopherol, beta-carotene, alpha-carotene, lycopene, and beta-cryptoxanthin in plasma, with tocopherol acetate as internal standard. Clinical Chemistry, 34, 377-381.
- Toskulkao, C. & Glinsukon, T. (1996): Endurance exercise and muscle damage: relationship to lipid peroxidation and scavenging enzymes in short and long distance runners. The Japanese Society of Physical Fitness and Sports Medicine, 45, 63-70.
- Urso, M.L. & Clarkson, P.M. (2003): Oxidative stress, exercise, and antioxidant supplementation. Toxicology, 15, 41-54.
- Visioli, F. & Galli, C. (2004): Olive oil: more than just oleic acid. American Journal of Clinical Nutrition, 72, 853.
- Vissers, M.N., Zock, P.L., Katan, M.B. (2004): Bioavailability and antioxidant effects of olive oil phenols in humans: a review. European Journal of Clinical Nutrition, 58, 955-965.
- Vissers, M.N., Zock, P.L., Leenen, R., Roodenburg, A.J.C., Van Putte, K.P.A.M. & Katan, M.B. (2001a): Effect of consumption of phenols from olives and extra virgin olive oil on LDL oxidizability in healthy humans. Free Radical Research, 35, 619-129.
- Vissers, M.N., Zock, P.L., Wiseman, S.A., Meyboom, S. & Katan, M.B. (2001b): Effect of phenol-rich extra virgin olive oil on markers of oxidation in healthy volunteers. European Journal of Clinical Nutrition, 55, 334-341.
- Vuilleumier, J.P.K. (1993): Fluorimetric assay of vitamin C in biological materials using a centrifugal analyser with fluorescence attachment. Journal of Micronutrient Analysis, 5, 25-34.
- Wiseman, S.A., Mathot, J.N., De Fouw, N. & Tijburg, L.B. (1996): Dietary non-tocopherol antioxidants present in extra virgin olive oil increase the resistance of low density lipoproteins to oxidation in rabbits. Atherosclerosis, 120, 15-23.

Submitted: 1-12-2009

Accepted: 11-2-2010

Correspondence to: I. Giannopoulou, MSc, PhD, Thorax Institute, School of Medicine, Athens, Greece,
Ploutarchou 3, Athens, fgiannopoulou@yahoo.gr

- References have been cited in the Greek (printed) version of the manuscript